



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A role for IL-15 in driving the onset of spontaneous autoimmune thyroiditis?

Citation for published version:

Kaiser, P, Rothwell, L, Vasícek, D & Hala, K 2002, 'A role for IL-15 in driving the onset of spontaneous autoimmune thyroiditis?', *The Journal of Immunology*, vol. 168, no. 8, pp. 4216-20.
<<http://www.jimmunol.org/content/168/8/4216.full.pdf+html>>

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

The Journal of Immunology

Publisher Rights Statement:

Copyright © 2002 by The American Association of Immunologists

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Take the Road Less Traveled.

**Bioactive Recombinant
Cytokines & Chemokines**

• Manufacturer of 170+ Proteins • Functional Testing on Every Lot



BioLegend®



A Role for IL-15 in Driving the Onset of Spontaneous Autoimmune Thyroiditis?

Pete Kaiser, Lisa Rothwell, Dusan Vasíček and Karel Hala

This information is current as of January 18, 2014.

J Immunol 2002; 168:4216-4220; ;
<http://www.jimmunol.org/content/168/8/4216>

-
- | | |
|----------------------|--|
| Subscriptions | Information about subscribing to <i>The Journal of Immunology</i> is online at:
http://jimmunol.org/subscriptions |
| Permissions | Submit copyright permission requests at:
http://www.aai.org/ji/copyright.html |
| Email Alerts | Receive free email-alerts when new articles cite this article. Sign up at:
http://jimmunol.org/cgi/alerts/etoc |

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
9650 Rockville Pike, Bethesda, MD 20814-3994.
Copyright © 2002 by The American Association of
Immunologists All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



A Role for IL-15 in Driving the Onset of Spontaneous Autoimmune Thyroiditis?¹

Pete Kaiser,^{2*} Lisa Rothwell,^{*} Dusan Vašiček,^{3†} and Karel Hala[†]

The obese strain (OS) of chickens, which suffers from spontaneous autoimmune thyroiditis, is an excellent animal model for Hashimoto's thyroiditis and provides a unique opportunity to investigate the mechanisms underlying and driving the onset of the disease. Following recent advances in cloning chicken cytokines, we can now begin to investigate the role of cytokines in driving the lymphoid infiltration of the thyroid seen in these birds from day 7 posthatch. Using real-time quantitative RT-PCR, we characterized the expression of IFN- γ , IL-1 β , IL-2, IL-6, IL-8, IL-15, and IL-18 in thyroids from OS birds and control CB line birds, both in the embryo just before hatch (embryonic day 20) and at 3 and 5 days posthatch. All of these cytokines were up-regulated compared with levels in thyroids from CB birds, at least at some time points, with some evidence for coordination of regulation, e.g., for the proinflammatory cytokines IL-1 β and IL-8. Only IL-15 was up-regulated at all time points. IL-15 was also shown to be up-regulated in spleens of OS birds at embryonic day 20 and 5 days posthatch, suggesting that IL-15 is constitutively up-regulated in this line of birds. This could explain the general immune system hyperreactivity exhibited by OS chickens and may be a factor driving the lymphoid infiltration of the thyroid. *The Journal of Immunology*, 2002, 168: 4216–4220.

Hashimoto's thyroiditis is a human organ-specific autoimmune disease that affects people of all ages, with peak occurrence in women of 30–50 years of age (1). Chickens of the obese strain (OS)⁴ with spontaneous autoimmune thyroiditis (SAT) have been used as an animal model for studies of this disease for the last 40 years (2, 3). All chickens from this strain develop symptoms of hypothyroidism due to SAT, and the immune system has been shown to be involved in the development of SAT. The onset of mononuclear cell infiltration of the thyroid in OS chicks is from 7 days of age, the first cells forming the multifocal infiltrates in the thyroid gland being activated CD4⁺, IL-2R⁺, TCR- $\alpha\beta_1$ ⁺ T cells (4). By 1 or 2 mo of age, the thyroid architecture is almost completely destroyed (5, 6). OS chickens exhibit general immune system hyperreactivity, believed in part to be due to intrinsic production of cytokines, perhaps IL-2 (7). Other immunomodulating factors in OS chickens that may contribute to SAT include premature emigration of Th cells from the thymus to the periphery during ontogenesis (8), an imbalance between effector and suppressor cells (9), and a disturbed immune-endocrine system interaction (6). Cytokines, in particular, probably have a role in initiating and perpetuating the disease.

Recent progress in the cloning of avian cytokines has led to the development of reagents with which to measure cytokine production in response to infection in the chicken. This should allow a greater insight into the mechanisms controlling the responses of the chicken to disease, both infectious and autoimmune, at both a cellular and molecular level. The avian orthologues of the Th1 cytokines IFN- γ , IL-2, and IL-18 recently have been cloned (10–12), as have the proinflammatory cytokines IL-1 β (13) and IL-6 (14); IL-15 (accession no. AF152927), which is closely related to IL-2; and the chemokine IL-8 (15, 16). The genomic sequences and gene structure for IFN- γ (17), IL-2 (18), IL-18 (P. Kaiser, unpublished data), IL-1 β (P. Kaiser, unpublished data), IL-15 (P. Kaiser, unpublished data), and IL-8 (19) have been fully determined. A partial genomic sequence for IL-6 has also been isolated recently (P. Kaiser, unpublished data). Gene structure information makes possible the design of probes and primers to specifically quantify cytokine mRNA levels using real-time quantitative RT-PCR.

We aimed to determine the levels of mRNA of these cytokines in the thyroids of 20-day-old embryos, 3- and 5-day-old OS, and unaffected CB birds to try to determine which cytokines, if any, might be driving the initial lymphocyte infiltration of the thyroid from 7 days of age in OS birds.

Materials and Methods

Experimental animals

OS leukosis-free chickens (5), bred as a closed flock homozygous for the MHC haplotype B13, were maintained under standardized conventional conditions (20) in the Central Laboratory Animal Facilities of the Medical Faculty, University of Innsbruck. Chickens of the unrelated leukosis-free inbred CB line (MHC haplotype B12) were used as a control line.

The designation of MHC haplotypes conforms to the nomenclature adopted at the International Workshop of the Chicken MHC (21). For details on the lines used, see the review by Hala and Plachy (22). To eliminate breeding errors, the MHC status of all animals was serologically determined.

Real-time quantitative RT-PCR

Cytokine mRNA levels in thyroids and spleens from OS and CB birds were quantified using a method based on that of Kaiser et al. (23).

Total RNA was prepared from pooled thyroids and spleens of five birds of each strain (OS and CB) at embryonic day 20 (E20), 3 days of age

*Institute for Animal Health, Compton, Berkshire, United Kingdom; and [†]Institute of Pathophysiology, University of Innsbruck, Innsbruck, Austria

Received for publication November 6, 2001. Accepted for publication February 15, 2002.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by Grant P12732-Med from the Fonds zur Förderung der Wissenschaftlichen Forschung and by the Biotechnology and Biological Sciences Research Council.

² Address correspondence and reprint requests to Dr. Pete Kaiser, Institute for Animal Health, Compton, Berkshire RG20 7NN, U.K. E-mail address: pete.kaiser@bbsrc.ac.uk

³ Current address: Research Institute of Animal Production, Institute of Genetics and Animal Reproduction, Hlohovská 2, 949 92 Nitra, Slovak Republic.

⁴ Abbreviations used in this paper: OS, obese strain; SAT, spontaneous autoimmune thyroiditis; E20, embryonic day 20; C_t, threshold cycle value; TAMRA, N,N,N,N'-tetramethyl-6-carboxyrhodamine.

Table I. Real-time quantitative RT-PCR probes and primers^a

RNA Target	Probe/Primer Sequence	Exon Boundary	Acc. No.
28S	Probe 5'-(VIC)-AGGACCGCTACGGACCTCCACCA-(TAMRA)-3' F 5'-GGCGAAGCCAGAGGAACT-3' R 5'-GACGACCGATTTCACGTC-3'		X59733
IFN- γ	Probe 5'-(FAM)-TGGCCAAGCTCCCGATGAACGA-(TAMRA)-3' F 5'-GTGAAGAAGGTGAAAGATATCATGGA-3' R 5'-GCTTTGCGCTGGATTCTCA-3'	3/4	Y07922
IL-1 β	Probe 5'-(FAM)-CCACACTGCAGCTGGAGGAAGCC-(TAMRA)-3' F 5'-GCTCTACATGTCGTGTGTGATGAG-3' R 5'-TGTCGATGTCCCGCATGA-3'	5/6	AJ245728
IL-2	Probe 5'-(FAM)-ACTGAGACCCAGGAGTGCACCCAGC-(TAMRA)-3' F 5'-TTGGAAAATATCAAGAACAAGATTCATC-3' R 5'-TCCCAGGTAACACTGCAGAGTTT-3'	2/3	AJ224516
IL-6	Probe 5'-(FAM)-AGGAGAAATGCCTGACGAAGCTCTCCA-(TAMRA)-3' F 5'-GCTCGCGGCTTCGA-3' R 5'-GGTAGGTCTGAAAGGCGAACAG-3'	3/4	AJ250838
IL-8	Probe 5'-(FAM)-TCTTTACCAGCGTCTACCTTGCGACA-(TAMRA)-3' F 5'-GCCCTCCTCCTGGTTTCA G-3' R 5'-TGGCACCGCAGCTCAT-3'	1/2	AJ009800
IL-15	Probe 5'-(FAM)-AAGTTGCAAATCTTGCAATTTCCATTTTTC-3' F 5'-TAGGAAGCATGATGTACGGAACAT-3' R 5'-TTTTTGCTGTTGTGGAATTCAACT-3'	4/5	AJ416937
IL-18	Probe 5'-(FAM)-CCGCGCCTTCAGCAGGGATG-(TAMRA)-3' F 5'-AGGTGAAATCTGGCAGTGGAAAT-3' R 5'-ACCTGGACGCTGAATGCAA-3'	4/5	AJ276026

^a Acc. No., genomic DNA sequence; F, forward; R, reverse; FAM, 5-carboxyfluorescein.

(thyroids only), and 5 days of age using Trizol reagent (Life Technologies, Gaithersburg, MD) following the manufacturer's instructions. Purified RNA was stored at -70°C .

For both cytokine and 28S rRNA-specific amplification, primers and probes were designed using the Primer Express software program (PE Applied Biosystems, Foster City, CA). Details of the probes and primers are given in Table I. All cytokine probes were designed, from the sequence of the relevant genes, to lie across intron:exon boundaries. Cytokine probes were labeled with the fluorescent reporter dye 5-carboxyfluorescein at the 5' end and with the quencher *N,N,N,N'*-tetramethyl-6-carboxyrhodamine (TAMRA) at the 3' end. The 28S probe was labeled with the fluorescent reporter dye VIC (PE Applied Biosystems) at the 5' end and with TAMRA at the 3' end.

RT-PCR was performed using the TaqMan EZ RT-PCR kit (PE Applied Biosystems). Amplification and detection of specific products were performed using the ABI PRISM 7700 Sequence Detection System (PE Applied Biosystems) with the following cycle profile: one cycle of 50°C for 2 min, 96°C for 5 min, 60°C for 30 min, and 95°C for 5 min, and 40 cycles of 94°C for 20 s, 59°C for 1 min.

Quantification was based on the increased fluorescence detected by the ABI PRISM 7700 Sequence Detection System due to hydrolysis of the target-specific probes by the 5' nuclease activity of the *rTth* DNA polymerase during PCR amplification. The passive reference dye 6-carboxy-x-rhodamine, which is not involved in amplification, was used to correct for fluorescent fluctuations, resulting from changes in the reaction conditions, for normalization of the reporter signal. Results are expressed in terms of the threshold cycle value (C_t), the cycle at which the change in the reporter dye passes a significance threshold. In this work, the threshold values of the change in the reporter dye are as shown in Table II for all reactions described.

To generate standard curves for the cytokine and 28S rRNA-specific reactions, total RNA, extracted from appropriately stimulated lymphoid cells, was serially diluted in sterile RNase-free water, and dilutions were made from 10^{-1} to 10^{-5} . Each RT-PCR experiment contained three no-template controls, test samples, and a \log_{10} dilution series. Each experiment was performed in triplicate, with replicates performed on different days. Regression analysis of the mean values of six replicate RT-PCRs for the \log_{10} diluted RNA was used to generate standard curves.

Results

Cytokine expression in the thyroid before the clinical onset of SAT

Replicate measurements on different days were highly repeatable, with a coefficient of variation for six replicate RT-PCRs of \log_{10}

serially diluted RNA for the different reactions, as shown in Table II. There was a linear relationship between the amount of input RNA and the C_t values for the various reactions as shown in Table II. Regression analyses of the C_t values generated by the \log_{10} dilution series gave R^2 values for all reactions in excess of 0.98 (see Table II for details). The increase in cycles per \log_{10} decrease in input RNA for each specific reaction, as calculated from the slope of the respective regression line, is given in Table II.

To control for variation in sampling and RNA preparation, the C_t values for cytokine-specific product for each sample were standardized using the C_t value of 28S rRNA product for the same sample. The C_t values for 28S rRNA did not alter significantly from sample to sample; the average 28S rRNA C_t values for all samples ranged from 9.12 to 10.46. Cytokine-specific C_t values varied from sample to sample and from cytokine to cytokine. The C_t values for 28S rRNA thus appeared to be independent of cytokine production and disease. Therefore, they were taken to be representative of the level of RNA extracted from all samples. To normalize RNA levels between samples within an experiment, the mean C_t value for 28S rRNA-specific product was calculated by pooling values from all samples in that experiment. Tube-to-tube variations in 28S rRNA C_t values about the experimental mean

Table II. Standard curve data from real-time quantitative RT-PCRs on total RNA extracted from stimulated splenocytes

	ΔRn^a	Log Dilutions	C_t	R^2	Slope
28S	0.05	10^{-1} – 10^{-5}	8–22	0.9833	3.0005
IFN- γ	0.01	10^{-1} – 10^{-5}	17–31	0.9899	3.289
IL-1 β	0.02	10^{-1} – 10^{-5}	24–38	0.9856	3.1553
IL-2	0.02	10^{-1} – 10^{-5}	26–38	0.9896	2.4279
IL-6	0.02	10^{-1} – 10^{-5}	23–37	0.9992	3.2841
IL-8	0.02	10^{-1} – 10^{-5}	14–27	0.9938	2.7728
IL-15	0.02	10^{-1} – 10^{-5}	22–36	0.9952	2.8154
IL-18	0.02	10^{-1} – 10^{-5}	17–33	0.9973	3.1123

^a ΔRn , Change in the reporter dye; C_t , the cycle at which the change in the reporter dye levels detected passes the ΔRn ; R^2 , coefficient of regression.




FIGURE 1. Quantification of IL-18 mRNA in RNA extracted from pooled thyroids from 20-day-old embryos and 3- and 5-day-old birds from either OS or CB strains (five birds per line per time point). C_t are expressed subtracted from 40 (the negative end point). Therefore, higher values represent higher levels of cytokine mRNA. □, Cytokine mRNA-specific C_t values before standardization for input RNA; ■, standardized values for cytokine mRNA corrected for variation in input RNA measured by 28S rRNA levels.

were calculated. The slope of the 28S rRNA \log_{10} dilution series regression line was used to calculate differences in input total RNA. Using the slopes of the respective cytokine \log_{10} dilution series regression lines, the difference in input total RNA, as represented by the 28S rRNA, was then used to adjust cytokine-specific C_t values. Fig. 1 shows the effect of standardizing cytokine-specific C_t values to correct for tube-to-tube variation in RNA levels. Standardization does not dramatically alter the distribution of the results as a whole.

IL-18 mRNA expression is up-regulated in OS E20 thyroids compared with CB E20 thyroids (Fig. 2). At 3 and 5 days posthatch, there is no difference in IL-18 mRNA expression between OS and CB thyroids. IFN- γ mRNA is down-regulated in OS E20 thyroids compared with CB E20 thyroids, it is of equal expression at 3 days posthatch, and it is up-regulated by 5 days posthatch (Fig. 2).

Up-regulation of the expression of the proinflammatory cytokines in OS thyroids compared with CB thyroids is biphasic (E20 and 5 days posthatch) for IL-1 β and IL-8, and at 5 days posthatch only for IL-6 (Fig. 3). At the other time points, there are no dif-




FIGURE 2. Quantification of IL-18 and IFN- γ mRNA in RNA extracted from pooled thyroids from 20-day-old embryos and 3- and 5-day-old birds from either OS or CB strains (five birds per line per time point), corrected for variation in input RNA as measured by 28S rRNA levels. The results are expressed as fold change in cytokine mRNA levels in pooled thyroids from OS birds when compared with those from age-matched CB birds (set as 1). No difference in mRNA levels, therefore, is shown as a fold change of 1. Error bars show SEM for triplicate samples from three separate TaqMan experiments.




FIGURE 3. Quantification of proinflammatory cytokine mRNA (IL-1 β , IL-6, and IL-8) in RNA extracted from pooled thyroids from 20-day-old embryos and 3- and 5-day-old birds from either OS or CB strains (five birds per line per time point), corrected for variation in input RNA as measured by 28S rRNA levels. The results are expressed as fold change in cytokine mRNA levels in pooled thyroids from OS birds when compared with those from age-matched CB birds (set as 1). No difference in mRNA levels, therefore, is shown as a fold change of 1. Error bars show SEM for triplicate samples from three separate TaqMan experiments.

ferences in mRNA expression between OS and CB thyroids for these cytokines, with the exception being IL-8 at 3 days posthatch, which is down-regulated in thyroids from OS birds.

IL-2 mRNA expression is up-regulated in OS thyroids compared with CB thyroids at 3 days posthatch only (Fig. 4). By contrast, IL-15 mRNA expression is up-regulated in OS thyroids at all time points in this study (Fig. 4).




FIGURE 4. Quantification of IL-15 and IL-2 mRNA in RNA extracted from pooled thyroids from 20-day-old embryos and 3- and 5-day-old birds from either OS or CB strains (five birds per line per time point), corrected for variation in input RNA as measured by 28S rRNA levels. The results are expressed as fold change in cytokine mRNA levels in pooled thyroids from OS birds when compared with those from age-matched CB birds (set as 1). No difference in mRNA levels, therefore, is shown as a fold change of 1. Error bars show SEM for triplicate samples from three separate TaqMan experiments.




FIGURE 5. Quantification of IL-15 mRNA in RNA extracted from pooled spleens (S) and thyroids (T) from 20-day-old embryos and 5-day-old birds from either OS or CB strains (five birds per line per time point), corrected for variation in input RNA as measured by 28S rRNA levels. The results are expressed as fold change in cytokine mRNA levels in pooled thyroids from OS birds when compared with those from age-matched CB birds (set as 1). Error bars show SEM for triplicate samples from three separate TaqMan experiments.

IL-15 mRNA expression is up-regulated in both spleens and thyroids of OS birds, both in the embryo and posthatch, compared with CB birds

Given the striking up-regulation of IL-15 mRNA in the thyroids of OS chickens at all stages sampled in this study compared with IL-15 mRNA levels in CB thyroids, we also investigated IL-15 mRNA expression in the spleens of E20 birds and birds 5 days posthatch. IL-15 mRNA expression is up-regulated in both the spleen and thyroid of OS birds at E20 and day 5 posthatch compared with CB birds (Fig. 5).

Discussion

IFN- γ has been shown to be up-regulated in the thyroids of patients suffering from Hashimoto's autoimmune thyroiditis (24–26) and in chickens after the onset of lymphoid infiltration of the thyroid in OS birds suffering from SAT (27). We have shown that IFN- γ expression in the embryonic thyroid before the onset of SAT in OS birds is down-regulated compared with CB chickens and that it is only up-regulated in the thyroid from OS birds 5 days posthatch, i.e., 2 days before the onset of lymphoid infiltration of the thyroid. In the thyroids of 5-day-old chicks, lymphocytes were not detected by immunohistochemistry (5). At this age, the thyroid mainly comprises cells of the thyroid follicular epithelium, fibroblasts, macrophages, and blood capillaries, none of which are able, under physiological conditions, to produce IFN- γ . Transgenic mice that constitutively express IFN- γ in the thyroid follicular cells developed severe hypothyroidism (28). In our model, the induction of expression of IFN- γ mRNA in the thyroid (Fig. 2), with subsequent lymphocyte infiltration, may be the result of a cytokine activation cascade initiated by IL-18 expression. This working concept of the mechanisms underlying the breakdown of self-tolerance and the induction of autoimmunity requires additional experiments.

IL-18 is not up-regulated in patients suffering from Hashimoto's thyroiditis (29), although such patients do have elevated IFN- γ levels (see above). In mammals, IL-18 has three main roles: to induce IFN- γ production (30), to enhance NK cell activity (31), and to activate neutrophils (32). To date, chicken IL-18 has only been shown to induce IFN- γ expression by splenocytes (12). It is interesting to note that thyroids from E20 OS birds have elevated IL-18 mRNA levels compared with age-matched thyroids from CB birds, suggesting that this elevated IL-18, if resulting in bioactive

protein, might be driving the elevated IFN- γ levels seen later in SAT.

Not surprisingly, considering the implication of T cell-mediated cytotoxic processes in the pathogenesis of Hashimoto's thyroiditis after the lymphoid infiltration of the thyroid, the proinflammatory cytokines IL-1 β , IL-6, and IL-8 have all been shown to be expressed in the thyroids of patients suffering from the disease (26, 33–36). This report shows that these cytokines are expressed in the thyroids of OS chickens before the onset of the lymphoid infiltration. Interestingly, for two of these cytokines (IL-1 β and IL-8) the up-regulation of their mRNA is biphasic, occurring both in the embryo (E20) and just before the onset of the lymphoid infiltration associated with the disease (day 5 posthatch). IL-6, in contrast, is only up-regulated 5 days posthatch.

IL-2 has been implicated in the pathogenesis of both Hashimoto's thyroiditis (33, 37) and SAT (7, 38). However, IL-2 mRNA is only up-regulated in the thyroids of OS birds at 3 days posthatch and therefore seems an unlikely candidate to be driving the lymphoid infiltration of the thyroid seen from day 7 posthatch. By contrast, the closely related cytokine IL-15 is up-regulated at all points in this study. In mammals, IL-15 shares many of the biological properties of IL-2. As well as being up-regulated in the thyroid of OS birds, IL-15 is also constitutively up-regulated in the spleens of OS birds at E20 and day 5 posthatch. Therefore, IL-15 represents a good candidate for the general immune system hyperactivity in OS birds (7) and may play a role in driving the lymphoid infiltration of the thyroid (IL-15 can be chemotactic for lymphocytes (39, 40)). There is one report of IL-15 mRNA being up-regulated in the thyroids of Hashimoto's thyroiditis patients (41).

At present, a lack of IL-15-specific reagents prevents us from investigating the role of this cytokine in SAT further. One obvious candidate for the overexpression of IL-15 in OS chickens would be a promoter polymorphism. However, although we have determined the gene structure for chicken IL-15, as yet we have no information on the promoter of chicken IL-15. Although bioactive recombinant chicken IL-15 has recently been expressed (42), as yet there are no mAbs described for chicken IL-15. However, mAbs specific for the α -chain of the IL-15R recently have been described (43), two of which inhibit the proliferative effect of IL-15 on T cells. This raises the possibility of blocking experiments in ovo and in vivo to determine whether IL-15 has a direct role in driving SAT, although these are far from simple in avian species because murine Abs do not fix chicken complement.

One potential drawback of real-time quantitative PCR is that, for cytokines, mRNA does not necessarily equate to bioactive protein. For example, both IL-1 β and IL-18 in mammals are produced initially as propeptides, which are then cleaved into an active form by the action of caspase-1 (44–47). The same mechanism is thought to apply for the avian orthologues of these two cytokines (12, 13). IL-15 production in mammals, in contrast, is predominantly controlled posttranslationally, and mRNA levels may be greater than protein levels (48–51). For other cytokines, mRNA levels correlate extremely well with bioactive protein. For example, in the chicken, IFN- γ and IL-6 mRNA levels (as measured by real-time quantitative PCR) and protein levels (as measured by bioassays) are in close agreement (23). However, we only have reliable bioassays for a limited number of chicken cytokines (type I IFN, IFN- γ , IL-1 β , IL-2, IL-6, and IL-18) and mAbs to even fewer (type I IFN, IFN- γ , and IL-2). Overall, real-time quantitative PCR is currently the only method that allows us to quantify a wide variety of avian cytokines within a particular disease model.

The question remains as to which cells in the thyroid of OS birds are expressing the cytokines we have identified in this study. To

this end, we intend to develop in situ hybridization for chicken cytokines in the thyroid, which in combination with immunohistochemistry should enable us to determine the phenotype of the cells expressing various cytokine mRNA before the onset of the lymphoid infiltration.

Acknowledgments

We thank Dr. K. Vašíčková and Dr. M. Kopečný for help with RNA isolation.

References

- Rose, N. R. 1991. The discovery of thyroid autoimmunity. *Immunol. Today* 12: 167.
- Sundick, R. S., N. Bagchi, and T. R. Brown. 1996. The obese strain chicken as a model for human Hashimoto's thyroiditis. *Exp. Clin. Endocrinol. Diabetes* 104(Suppl. 3):4.
- Wick, G., J. Moest, K. Schauenstein, G. Krömer, H. Dietrich, A. Ziemiecki, R. Fässler, S. Schwarz, N. Neu, and K. Hala. 1985. Spontaneous autoimmune thyroiditis: a bird's eye view. *Immunol. Today* 6:359.
- Cihak, J., G. Hoffman-Fezer, M. Wasl, H. Merkle, B. Kaspers, O. Vainio, J. Plachy, K. Hala, G. Wick, M. Stangassinger, and U. Lösch. 1998. Inhibition of the development of spontaneous autoimmune thyroiditis in the obese strain (OS) chickens by in vivo treatment with anti-CD4 or anti-CD8 Abs. *J. Autoimmun.* 11:119.
- Hala, K., G. Malin, H. Dietrich, U. Lösch, G. Böck, H. Wolf, B. Kaspers, J. Geryk, M. Falk, and R. L. Boyd. 1996. Analysis of the initiation period of spontaneous autoimmune thyroiditis (SAT) in obese strain (OS) of chickens. *J. Autoimmun.* 9:129.
- Wick, G., H. P. Brezinschek, K. Hala, H. Dietrich, H. Wolf, and G. Kroemer. 1989. The obese strain of chickens: an animal model with spontaneous autoimmune thyroiditis. *Adv. Immunol.* 47:433.
- Schauenstein, K., G. Kroemer, R. S. Sundick, and G. Wick. 1985. Enhanced response to Con A and production of TCGF by lymphocytes of obese strain (OS) chickens with spontaneous autoimmune thyroiditis. *J. Immunol.* 134:872.
- Rose, N. R., L. D. Bacon, and R. S. Sundick. 1976. Genetic determinants of thyroiditis in the OS chicken. *Transplant. Rev.* 31:264.
- Boyd, R. L., K. Hala, and G. Wick. 1985. Interactions and quantitative analysis of immunoregulatory cells in the chicken thymus. *J. Immunol.* 135:3039.
- Digby, M. R., and J. W. Lowenthal. 1995. Cloning and expression of the chicken interferon- γ gene. *J. Interferon Cytokine Res.* 15:939.
- Sundick, R. S., and C. Gill-Dixon. 1997. A cloned chicken lymphokine homologous to both mammalian IL-2 and IL-15. *J. Immunol.* 159:720.
- Schneider, K., F. Puehler, D. Baeuerle, S. Elvers, P. Staeheli, B. Kaspers, and K. C. Weining. 2000. cDNA cloning of biologically active chicken interleukin-18. *J. Interferon Cytokine Res.* 20:879.
- Weining, K. C., C. Sick, B. Kaspers, and P. Staeheli. 1998. A chicken homolog of mammalian interleukin-1 β : cDNA cloning and purification of active recombinant protein. *Eur. J. Biochem.* 258:994.
- Schneider, K., R. Klaas, B. Kaspers, and P. Staeheli. 2001. Chicken interleukin-6 cDNA structure and biological properties. *Eur. J. Biochem.* 268:4200.
- Bedard, P.-A., D. Alcorta, D. L. Simmons, K.-C. Luk, and R. L. Erikson. 1987. Constitutive expression of a gene encoding a polypeptide homologous to biologically active human platelet protein in Rous sarcoma virus-transformed fibroblasts. *Proc. Natl. Acad. Sci. USA* 84:6715.
- Sugano, S., M. Y. Stoeckle, and H. Hanafusa. 1987. Transformation by Rous sarcoma virus induces a novel gene with homology to a mitogenic platelet protein. *Cell* 49:321.
- Kaiser, P., H. M. Wain, and L. Rothwell. 1998. Structure of the chicken interferon- γ gene, and comparison to mammalian homologues. *Gene* 207:25.
- Kaiser, P., and P. Mariani. 1999. Promoter sequence, exon:intron structure and synteny of genetic location show that a chicken cytokine with T cell proliferative activity is IL2 and not IL15. *Immunogenetics* 49:26.
- Kaiser, P., S. Hughes, and N. Bumstead. 1999. The chicken 9E3/CEF4 CXC chemokine is the avian orthologue of IL8 and maps to chicken chromosome 4 syntenic with genes flanking the mammalian chemokine cluster. *Immunogenetics* 49:673.
- Dietrich, H. M. 1989. Housing, breeding and selecting chickens of the Obese strain (OS) with spontaneous autoimmune thyroiditis. *Lab. Anim.* 23:345.
- Briles, W. E., N. Bumstead, D. L. Ewert, D. G. Gilmour, J. Gogusev, K. Hala, C. Koch, B. M. Longenecker, A. W. Nordskog, J. R. Pink, et al. 1982. Nomenclature for chicken major histocompatibility (B) complex. *Immunogenetics* 5:441.
- Hala, K., and J. Plachy. 1997. Inbred strain of chickens. In *Immunology Methods Manual*, I. Lefkowitz, ed. Academic Press, New York, p. 2285.
- Kaiser, P., L. Rothwell, E. E. Galyov, P. A. Barrow, J. Burnside, and P. Wigley. 2000. Differential cytokine expression in avian cells in response to invasion by *Salmonella typhimurium*, *Salmonella enteritidis* and *Salmonella gallinarum*. *Microbiology* 146:3217.
- Bagnasco, M., S. Ferrini, D. Venuti, I. Prigione, G. Torre, R. Biassoni, and G. W. Canonica. 1987. Clonal analysis of T lymphocytes infiltrating the thyroid gland in Hashimoto's thyroiditis. *Int. Arch. Allergy Appl. Immunol.* 82:141.
- Del Prete, G. F., A. Tiri, S. Mariotti, A. Pinchera, M. Ricci, and S. Romagnani. 1987. Enhanced production of γ -interferon by thyroid-derived T cell clones from patients with Hashimoto's thyroiditis. *Clin. Exp. Immunol.* 69:323.
- Ajjan, R. A., P. F. Watson, R. S. McIntosh, and A. P. Weetman. 1996. Intrathyroidal cytokine gene expression in Hashimoto's thyroiditis. *Clin. Exp. Immunol.* 105:523.
- Hempel, K., P. Schmidt, W. Schmahl, K. Hala, H.-G. Wick, and B. Kaspers. 2001. The role of interferon- γ and the inducible nitric oxide synthase in the pathophysiology of the spontaneous autoimmune thyroiditis of the Obese strain of chicken. In *Current Progress on Avian Immunology Research*, K. A. Schat, ed. American Association of Avian Pathologists, Kennett Square, PA, p. 144.
- Caturegli, P., M. Hejazi, K. Suzuki, O. Dohan, N. Carrasco, L. D. Kohn, and N. R. Rose. 2000. Hypothyroidism in transgenic mice expressing IFN- γ in the thyroid. *Proc. Natl. Acad. Sci. USA* 97:1719.
- Miyauchi, S., B. Matsuura, and M. Onji. 2000. Increased levels of serum interleukin-18 in Graves' disease. *Thyroid* 10:815.
- Okamura, H., H. Tsutsui, T. Komatsu, M. Yutsudo, A. Hakura, T. Tanimoto, K. Torigoe, T. Okura, Y. Nukada, K. Hattori, et al. 1995. Cloning of a new cytokine that induces IFN- γ production by T cells. *Nature* 378:88.
- Tsutsui, H., K. Nakanishi, K. Matsui, K. Higashino, H. Okamura, Y. Miyazawa, and K. Kaneda. 1996. IFN- γ -inducing factor up-regulates Fas ligand-mediated cytotoxic activity of murine natural killer cell clones. *J. Immunol.* 157:3967.
- Leung, B. P., S. Culshaw, J. A. Gracie, D. Hunter, C. A. Canetti, C. Campbell, F. Cunha, F. Y. Liew, and I. B. McInnes. 2001. A role for IL-18 in neutrophil activation. *J. Immunol.* 167:2879.
- Paschke, R., F. Schuppert, M. Taton, and T. Velu. 1994. Intrathyroidal cytokine gene expression profiles in autoimmune thyroiditis. *J. Endocrinol.* 141:309.
- Kayser, L., H. Broholm, D. Francis, H. Perrild, B. E. Olsen, K. Bendtzen, and P. E. Hoyer. 1995. Immunocytochemical localisation of interleukin-1 α and interleukin-6 in thyroid tissues from patients with neoplastic or autoimmune thyroid disorders. *Autoimmunity* 20:75.
- Heuer, M., G. Aust, S. Ode-Hakim, and W. A. Scherbaum. 1996. Different cytokine mRNA profiles in Graves' disease, Hashimoto's thyroiditis, and nonautoimmune thyroid disorders determined by quantitative reverse transcriptase polymerase chain reaction (RT-PCR). *Thyroid* 6:97.
- Paolieri, F., C. Salmaso, M. Battifora, P. Montagna, G. Pesce, M. Bagnasco, R. Richiusa, A. Galluzzo, and C. Giordano. 1999. Possible pathogenetic relevance of interleukin-1 β in "destructive" organ-specific autoimmune disease (Hashimoto's thyroiditis). *Ann. NY Acad. Sci.* 876:221.
- Drugarin, D., S. Negru, A. Koreck, I. Zosin, and C. Cristea. 2000. The pattern of a T(H)1 cytokine in autoimmune thyroiditis. *Immunol. Lett.* 71:73.
- Kromer, G., R. S. Sundick, K. Schauenstein, K. Hala, and G. Wick. 1985. Analysis of lymphocytes infiltrating the thyroid gland of Obese strain chickens. *J. Immunol.* 135:2452.
- Jonuleit, H., K. Wiedemann, G. Muller, J. Degwert, U. Hoppe, J. Knop, and A. H. Henk. 1997. Induction of IL-15 messenger RNA and protein in human blood-derived dendritic cells: a role for IL-15 in attraction of T cells. *J. Immunol.* 158:2610.
- Sancho, D., M. Yanez-Mo, R. Tejedor, and F. Sanchez-Madrid. 1999. Activation of peripheral blood T cells by interaction and migration through endothelium: role of lymphocyte function antigen-1/intercellular adhesion molecule-1 and interleukin-15. *Blood* 93:886.
- Ajjan, R. A., P. F. Watson, and A. P. Weetman. 1997. Detection of IL-12, IL-13, and IL-15 messenger ribonucleic acid in the thyroid of patients with autoimmune thyroid disease. *J. Clin. Endocrinol. Metab.* 82:666.
- Lillehoj, H. S., W. Min, K. D. Choi, U. S. Babu, J. Burnside, T. Miyamoto, B. M. Rosenthal, and E. P. Lillehoj. 2001. Molecular, cellular, and functional characterization of chicken cytokines homologous to mammalian IL-15 and IL-2. *Vet. Immunol. Immunopathol.* 82:229.
- Li, G., H. S. Lillehoj, and W. Min. 2001. Production and characterisation of monoclonal Abs reactive with the chicken interleukin-15 receptor α chain. *Vet. Immunol. Immunopathol.* 82:215.
- Black, R. A., S. R. Kronheim, J. E. Merriam, C. J. March, and T. P. Hopp. 1989. A pre-aspartate-specific protease from human leukocytes that cleaves pro-interleukin-1 β . *J. Biol. Chem.* 264:5323.
- Kostura, M. J., M. J. Tocci, G. Limjuco, J. Chin, P. Cameron, A. G. Hillman, N. A. Chartrain, and J. A. Schmidt. 1989. Identification of a monocyte specific pre-interleukin 1 β convertase activity. *Proc. Natl. Acad. Sci. USA* 86:5227.
- Ghayur, T., S. Banerjee, M. Hugunin, D. Butler, L. Herzog, A. Carter, L. Quintal, L. Sekut, R. Talanian, M. Paskind, et al. 1997. Caspase-1 processes IFN- γ -inducing factor and regulates LPS-induced IFN- γ production. *Nature* 386:619.
- Gu, Y., K. Kuida, H. Tsutsui, G. Ku, K. Hsiao, M. A. Fleming, N. Hayashi, K. Higashino, H. Okamura, K. Nakanishi, et al. 1997. Activation of interferon- γ inducing factor mediated by interleukin-1 β converting enzyme. *Science* 275:206.
- Grabstein, K. H., J. Eisenman, K. Shanebeck, C. Rauch, S. Srinivasan, V. Fung, C. Beers, J. Richardson, M. A. Schoenborn, M. Ahdieh, et al. 1994. Cloning of a T cell growth factor that interacts with the β chain of the interleukin-2 receptor. *EMBO J.* 14:3654.
- Bamford, R. N., A. P. Battista, J. D. Burton, H. Sharma, and T. A. Waldmann. 1996. Interleukin (IL) 15/IL-T production by the adult T-cell leukemia cell line HuT-102 is associated with a human T-cell lymphotropic virus type II region/IL-15 fusion message that lacks many upstream AUGs that normally attenuate IL-15 mRNA translation. *Proc. Natl. Acad. Sci. USA* 93:2897.
- Tagaya, Y., R. N. Bamford, A. P. de Felippis, and T. A. Waldmann. 1996. IL-15: a pleiotropic cytokine with diverse receptor/signaling pathways whose expression is controlled at multiple levels. *Immunity* 4:329.
- Tagaya, Y., G. Kurys, T. A. Thies, J. M. Losi, N. Azimi, J. A. Hanover, R. N. Bamford, and T. A. Waldmann. 1997. Generation of secreted and non-secreted interleukin 15 isoforms through alternate usage of signal peptides. *Proc. Natl. Acad. Sci. USA* 94:14444.